

# Tuberculosis (TB) Vaccine Research Study



## International Travelers are needed for a BCG TB vaccine study

- **Healthcare Worker (HCW) traveling**  
≥ 4 weeks, but < 6 months,

**OR**

- **HCW/non-HCW traveling**  
≥ 6 months to 30 months (2 and a half years)

### BCG TIPI STUDY

(See [www.clinicaltrials.gov](http://www.clinicaltrials.gov) Search Identifier: NCT04453293)

Researchers are evaluating whether pre-travel investigational BCG immunization can prevent TB infection in an “at-risk” traveler population traveling to countries with a high burden of TB where there is a risk for multidrug resistant and extensively drug-resistant TB exposure. Information collected may provide valuable information to further describe and identify TB exposure/risk factors for infection as well as any information regarding development of active TB disease, and the efficacy of BCG vaccine to prevent symptomatic all-cause respiratory infections during travel abroad.

#### You may be eligible if:

- ◆ Between 18 and 65 years of age
- ◆ In good health and HIV negative
- ◆ Have not received BCG vaccine
- ◆ Never had TB infection before
- ◆ Planning travel to country with high TB rate
- ◆ Not pregnant or lactating

#### Study procedures include:

- ◆ In-person visits (4-6) at the study site
- ◆ General health screening
- ◆ Blood draws (2 up to 4)
- ◆ BCG vaccination or placebo
- ◆ Study diary & questionnaires completion
- ◆ Compensation will be provided for study visits completed (about \$150/completed study visit)

**Study Site Location: 855 N Wolfe St Rm 601, Baltimore MD, 21205**  
**If interested or would like more information, please call: 410-955-7283**