

Tuberculosis (TB) Vaccine Research Study



International Travelers needed for a BCG TB vaccine study

- **Healthcare or Public Health workers traveling to a TB-endemic country for a minimum of 10 days and no longer than 6 months**

BCG TIPI STUDY

(See www.clinicaltrials.gov Search Identifier: NCT04453293)

Researchers are evaluating whether pre-travel investigational BCG immunization can prevent infections in an “at-risk” traveler population traveling to countries with a moderate and high burden of TB where there is a risk for multidrug resistant and extensively drug-resistant TB exposure. Information collected may provide valuable information to further describe and identify exposure/risk factors for infection as well as any information regarding development of active TB disease, and the efficacy of BCG vaccine to prevent symptomatic all-cause respiratory infections during travel abroad.

You may be eligible if:

- ◆ Between 18 and 65 years of age
- ◆ In good health and HIV negative
- ◆ Have not received BCG vaccine
- ◆ Never had TB infection before
- ◆ Planning travel to country with high - moderate TB rate
- ◆ Not pregnant or lactating

Study procedures include:

- ◆ In-person visits (4-6) at the study site
- ◆ General health screening
- ◆ Blood draws (2 up to 4)
- ◆ BCG vaccination or placebo
- ◆ Study diary & questionnaires completion
- ◆ Compensation will be provided for study visits completed (about \$150/completed study visit)

Study Site Location: 855 N Wolfe St Suite 601, Baltimore MD, 21205

If interested or would like more information, please contact: TIPI@jh.edu, call 336-660-9392, or visit <https://centerforimmunizationresearch.org>.